arXiv:1703.00668v1 [math.DS] 2 Mar 2017

PERIODIC POINTS OF ALGEBRAIC ACTIONS OF DISCRETE GROUPS

SIDDHARTHA BHATTACHARYA

Abstract. Let Γ be a countable group. A Γ-action on a compact abelian group X by continuous automorphisms of X is called Noetherian if the dual of X is Noetherian as a $\mathbb{Z}(\Gamma)$-module. We prove that any Noetherian action of a finitely generated virtually nilpotent group has a dense set of periodic points.

1. Introduction

Let Γ be a countable group. An *algebraic Γ-action* is an action α of Γ on a compact metrizable abelian group X by continuous automorphisms of X. By duality theory, any such action induces a Γ-action on $\widehat{X}$, the dual of X, by automorphisms of $\widehat{X}$. Hence $\widehat{X}$ can be viewed as a $\mathbb{Z}[\Gamma]$-module, where $\mathbb{Z}[\Gamma]$ is the integral group ring of Γ. The action α is called *Noetherian* if $\widehat{X}$ is Noetherian as a $\mathbb{Z}[\Gamma]$-module. Equivalently, α is Noetherian if any decreasing sequence

$$X = X_0 \supset X_1 \supset X_2 \supset \cdots$$

of closed Γ-invariant subgroups stabilizes.

For $\Gamma = \mathbb{Z}^d$, the study of such actions was initiated in [5], and since then these systems have been extensively studied (see [9] for a comprehensive account). For general Γ, several important results about entropy and homoclinic points of Noetherian Γ-actions have been proved in recent years ([1], [2], [3], [6]). However, unlike the commutative case, many basic dynamical properties are only partly understood in the general situation.

In this paper, we study periodic points of Noetherian Γ-actions. Recall that a point $x \in X$ is periodic if the Γ-orbit of x is finite. It is

2010 *Mathematics Subject Classification.* 37B05, 37B20.

Key words and phrases. Nilpotent groups, automorphisms, periodic points.

known that when $\Gamma = \mathbb{Z}^d$, for any Noetherian Γ-action (X, α) the space X contains a dense set of periodic points ([9], Theorem 5.7). However, the proof uses tools from commutative algebra and does not generalize to actions of non-abelian groups. The question whether Noetherian actions of all residually finite groups admit a dense set of periodic points was raised in [7, Problem 8.5].

In this paper we show that for algebraic actions of countable groups, the density of periodic points is related to vanishing of certain cohomology groups. As a consequence, we obtain the following :

Theorem 1.1. *Let Γ be a finitely generated virtually nilpotent group and let α be a Noetherian action of Γ on a compact metrizable abelian group X. Then X contains a dense set of α-periodic points.*

As another application we show that there exist Noetherian actions of finitely generated residually finite groups that do not have a dense set of periodic points, thus giving a negative answer to the above question in the most general case.

2. Virtual first cohomology groups

Let α be an action of a countable group Γ on a compact metrizable group X by continuous automorphisms of X. A *virtual 1-cocycle* of α is a map c from a finite index subgroup $\Lambda \subset \Gamma$ to X that satisfies the equation

$$c(\gamma\gamma^{'}) = c(\gamma) + \alpha(\gamma)(c(\gamma^{'}))$$

for all $\gamma, \gamma^{'} \in \Lambda$. Two virtual 1-cocycles $c_1 : \Lambda_1 \to X$ and $c_2 : \Lambda_2 \to X$ are *equivalent* if there exists a finite index subgroup $\Lambda \subset \Lambda_1 \cap \Lambda_2$ such that $c_1(\gamma) = c_2(\gamma)$ for all $\gamma \in \Lambda$. For any $x \in X$, the map $c_x : \Gamma \to X$ defined by $c_x(\gamma) = \alpha(\gamma)(x) - x$ is a 1-cocycle. A virtual 1-cocycle $c : \Lambda \to X$ is said to be a *virtual coboundary* if it is equivalent to c_x for some $x \in X$. We will call two virtual 1-cocycles $c_1 : \Lambda_1 \to X$ and $c_2 : \Lambda_2 \to X$ *cohomologous* if there exists a finite index subgroup $\Lambda \subset \Lambda_1 \cap \Lambda_2$, and a virtual coboundary $c : \Lambda \to X$ such that $c_1(\gamma) - c_2(\gamma) = c(\gamma)$ for all $\gamma \in \Lambda$. It is easy to see that the equivalence classes of virtual 1-cocyles is a group with respect to the pointwise addition, and the equivalence classes of virtual coboundaries form a subgroup.

We will denote the quotient group by $H^1_v(\alpha)$ and call it the *virtual first cohomology group* of α.

Let (Y, β) be an algebraic Γ-action, and let $X \subset Y$ be a closed β-invariant subgroup such that Y/X is finite. Let α denote the restriction of β to X and let $\Lambda \subset \Gamma$ be a finite index subgroup that acts trivially on Y/X. For any $y \in Y$ we define a map $c_y : \Lambda \to X$ by $c_y(\gamma) = \beta(\gamma)(y) - y$. It is easy to see that c_y is a virtual 1-cocycle of α.

Proposition 2.1. *Let (Y, β), (X, α) and $y \in Y$ be as above. Then the virtual 1-cocycle c_y is a virtual coboundary if and only if the coset $y + X$ contains a β-periodic point.*

Proof. Suppose $y_1 \in y + X$ is a β-periodic point, and $\Lambda_1 \subset \Lambda$ is a finite index subgroup such that $\beta(\gamma)(y_1) = y_1$ for all $\gamma \in \Lambda_1$. Then $x = y - y_1 \in X$ and

$$c_y(\gamma) = \beta(\gamma)(y) - y = \beta(\gamma)(x) - x$$

for all $\gamma \in \Lambda_1$. Hence c_y is a virtual coboundary of α. Conversely, if c_y is a virtual coboundary of α, then there exists a finite index subgroup Λ_1 and $x \in X$ such that $c_y(\gamma) = \beta(\gamma)(x) - x$ for all $\gamma \in \Lambda_1$. If $p = y - x$ then $p \in y + X$ and it is fixed by $\beta(\Lambda_1)$. Hence it is a periodic point. □

A compact abelian group X is *zero-dimensional* if the connected component containing 0_X is trivial. From the duality theory it follows that X is zero-dimensional if and only if every element of $\widehat{X}$ has finite order.

Proposition 2.2. *Let Γ be a countable group, and let α be a Noetherian action of Γ on a zero-dimensional compact abelian group X. Then there exists $k \geq 1$ such that $kx = 0$ for all $x \in X$.*

Proof. Let $\{\chi_1, \ldots, \chi_n\} \subset \widehat{X}$ be a finite set that generates $\widehat{X}$ as a $\mathbb{Z}(\Gamma)$-module. As every element of $\widehat{X}$ has finite order, we can find $k \geq 1$ such that $k\chi_i = 0$ for $i = 1, \ldots, n$. Since for any m the set $N_m = \{\chi : m\chi = 0\}$ is a $\mathbb{Z}(\Gamma)$-submodule of $\widehat{X}$, we deduce that $N_k = \widehat{X}$, i.e., $k\chi = 0$ for all $\chi \in \widehat{X}$. If $x \in X$ then for all $\chi \in \widehat{X}$, $\chi(kx) = k\chi(x) = 0$. Hence $kx = 0$. □

Our next proposition shows that all virtual 1-cocycles of Noetherian Γ-actions on zero-dimensional groups arise in the manner described in Proposition 2.1.

Proposition 2.3. *Let Γ be a countable group and let (X, α) be a Noetherian action of Γ on a zero-dimensional group X. Then for any virtual 1-cocycle $c : \Lambda \to X$ there exists $k \geq 1$ and an algebraic Λ-action β on $Y = (\mathbb{Z}/k\mathbb{Z}) \times X$ such that*

(1) *For all $\gamma \in \Lambda$, $\beta(\gamma)$ fixes the first co-ordinate.*
(2) *$\beta(\gamma)(0, x) = (0, \alpha(\gamma)(x))$ for all $\gamma \in \Lambda$ and $x \in X$.*
(3) *$c_{(1,0)} = c$.*

Proof. Since α is Noetherian and X is zero-dimensional, by the previous proposition there exists $k \geq 1$ such that $kx = 0$ for all $x \in X$. For any $z \in Y$ and a continuous automorphism τ of X there exists a unique homomorphism $\theta : Y \to Y$ such that $\theta(1, 0) = z$ and $\theta(0, x) = (0, \tau(x))$ for all $x \in X$. It is easy to see that θ is continuous. For $\gamma \in \Lambda$ let $\beta(\gamma)$ denote the unique continuous endomorphism of Y satisfying the following two conditions :

(1) $\beta(\gamma)(0, x) = (0, \alpha(\gamma)(x))$ for all $x \in X$.
(2) $\beta(\gamma)(1, 0) = (1, c(\gamma))$.

Similarly we define $\beta'(\gamma)$ to be the unique continuous endomorphism of Y with the property that $\beta'(\gamma)(0, x) = (0, \alpha(\gamma)^{-1}(x))$ and $\beta'(\gamma)(1, 0) = (1, -\alpha(\gamma)^{-1}c(\gamma))$. Then

$$\beta(\gamma)\beta'(\gamma)(1, 0) = \beta(\gamma)(1, 0) + \beta(\gamma)(0, -\alpha(\gamma)^{-1}c(\gamma)) = (1, 0).$$

As $\beta(\gamma)\beta'(\gamma)$ fixes both $\{0\} \times X$ and the point $(1, 0)$, we conclude that $\beta(\gamma)\beta'(\gamma) = I$. A similar computation shows that $\beta'(\gamma)\beta(\gamma) = I$. Hence $\beta(\gamma)$ is a continuous automorphism of Y for each $\gamma \in \Lambda$. As $c(\gamma_1\gamma_2) = c(\gamma_1) + \alpha(\gamma_1)c(\gamma_2)$, it follows that

$$\beta(\gamma_1\gamma_2)(1, 0) = (1, c(\gamma_1\gamma_2)) = \beta(\gamma_1)\beta(\gamma_2)(1, 0).$$

Since $\beta(\gamma_1\gamma_2)(0, x) = \beta(\gamma_1)\beta(\gamma_2)(0, x)$ for all $x \in X$, this imlies that $\beta(\gamma_1\gamma_2) = \beta(\gamma_1)\beta(\gamma_2)$. Hence β defines an algebraic Λ-action on Y. It is easy to see that β satisfies all three conditions stated in the proposition. □

Lemma 2.4. *Let Γ be a countable group and let (X, α) be a Noetherian Γ-action such that X admits non-zero periodic points. Then X also contains non-trivial α-invariant finite subgroups.*

Proof. Let $x \in X$ be a non-zero α-periodic point and let $\Lambda \subset \Gamma$ be the stabilizer of x. Since the orbit of x is finite, Λ is a finite index subgroup of Γ. We define a subgroup $\Lambda_0 \subset \Gamma$ by

$$\Lambda_0 = \bigcap_{\gamma \in \Gamma} \gamma \Lambda \gamma^{-1}.$$

It is easy to see that $\Lambda_0 \subset \Gamma$ is normal and has finite index. Let Y denote the set of points of X that are fixed by all elements of Λ_0. Then Y is a non-trivial α-invariant closed subgroup of X. Since $\Lambda_0 \subset \Gamma$ has finite index, the action $\alpha|_{\Lambda_0}$ is also Noetherian. As Λ_0 acts trivially on Y, it follows that any collection of closed subgroups of Y has a minimal element. Let Y_0 be a minimal element of the collection of all non-trivial subgroups of Y. Since closed subgroups of Y_0 are in one to one correspondence with subgroups of $\widehat{Y_0}$, we deduce that $\widehat{Y_0}$ does not admit non-trivial subgroups. Therefore $\widehat{Y_0}$, and hence Y_0, are isomorphic with $\mathbb{Z}/p\mathbb{Z}$ for some prime p. In particular, Y_0 is finite. We define $H \subset X$ by

$$H = \sum_{\gamma \in \Gamma} \alpha(\gamma)(Y_0).$$

As $\alpha(\gamma)(Y_0) = Y_0$ for all $\gamma \in \Lambda_0$, and $\Lambda_0 \subset \Gamma$ has finite index, the above sum is finite and H is a well defined closed subgroup of X. It is easy to see that H is finite and α-invariant. $\square$

The above lemma is not true if α is not assumed to be Noetherian. For example, suppose Γ is an arbitrary countable group and α is the trivial action of Γ on $X = \widehat{\mathbb{Q}}$. Then every point of X is α-periodic, but since the group $\mathbb{Q}$ does not have non-trivial finite quotients, X does not admit non-trivial finite subgroups.

Theorem 2.5. *Let Γ be a countable group satisfying :*

1. *Every Noetherian Γ-action (X, α) with $X \neq \{0\}$, has a non-zero α-periodic point.*
2. *The group $H^1_v(\alpha)$ is trivial for every Noetherian Γ-action (X, α) on a zero-dimensional group X.*

Then every Noetherian action of Γ *on a compact abelian group (not necessarily zero-dimensional) admits a dense set of periodic points. Conversely, if every Noetherian* Γ*-action admits a dense set of periodic points, then both the conditions are satisfied.*

Proof. Suppose every Noetherian action of Γ admits a dense set of periodic orbits. Then the first condition is automatically satisfied. Let α be a Noetherian Γ-action on a zero-dimensional group X, and let $c : \Lambda \to X$ be a virtual 1-cocycle of α. By Proposition 2.3, there exists $k \geq 1$ and an algebraic Λ-action β on $Y = (\mathbb{Z}/k\mathbb{Z}) \times X$ such that $c = c_{(1,0)}$ and $\beta(\gamma)|_X = \alpha(\gamma)$ for all $\gamma \in \Lambda$. Since the coset containing $(1, 0)$ is an open subset of Y, it contains a β-periodic orbit. From Proposition 2.1 we deduce that c is a virtual coboundary. Hence $H^1_v(\alpha) = \{0\}$.

Now suppose Γ satisfies both the conditions stated above. Let (Y, β) be a Noetherian action of Γ. Let $X \subset Y$ denote the closure of the set of all β-periodic points. Since the sum of two periodic points is again a periodic point it follows that X is a closed β-invariant subgroup. Suppose $X \neq Y$. Let $\bar{\beta}$ denote the induced action of Γ on the quotient Y/X. Since Γ satisfies the first condition, by Lemma 2.4 there exists a finite $\bar{\beta}$-invariant non-zero subgroup $F \subset Y/X$. We define $Y' = \pi^{-1}(F)$, where π is the projection map from Y to Y/X. We choose a finite index subgroup $\Lambda \subset \Gamma$ that acts trivially on F under $\bar{\beta}$. Let $y \notin X$ be a point in Y', and let $c_y : \Lambda \to X$ denote the virtual 1-cocycle of (X, β) defined by $c_y(\gamma) = \beta(\gamma)(y) - y$.

Let X_0 denote the connected component of X. It is easy to see that X_0 is a β-invariant closed subgroup of X. Let β_1 denote the induced action of Λ on X/X_0. As X/X_0 is zero-dimensional and β_1 is Noetherian, applying Proposition 2.2 we deduce that there exists $l \geq 2$ such that $lp \in X_0$ for all $p \in X$. Let $k \geq 2$ be a positive integer such that $kx = 0$ for all $x \in F$. Then $lky \in X_0$. Since X_0 is connected, the map $x \mapsto lkx$ is a surjective endomorphism of X_0. We find $q \in X_0$ such that $lky = lkq$, and define a virtual 1-cocycle $c_1 : \Lambda \to X$ by $c_1 = c_y - c_q$. Then for any $\gamma \in \Lambda$,

$$lkc_1(\gamma) = lk(\beta(\gamma)(y - q) - (y - q)) = 0.$$

Hence the image of c_1 is contained in $H = \{x \in X : lkx = 0\}$. Since H is closed, β-invariant, and zero-dimensional; from the second condition we conclude that c_1 is a virtual coboundary. As c_1 is cohomologous to c_y, this implies that c_y is also a virtual coboundary. Applying Proposition 2.1 we deduce that the coset $y + X$ contains a periodic point y_1. This contradicts the fact that $y \notin X$, and shows that $Y = X$, i.e., the set of β-periodic points is dense in Y. $\square$

We now construct an example of a Noetherian action of a finitely generated residually finite group that does not have a dense set of periodic points. Let H denote the group of all functions from $\mathbb{Z}$ to $\mathbb{Z}$ with finite support, equipped with pointwise addition. We define an action of $\mathbb{Z}$ on H by $n \cdot f(i) = f(i + n)$. Let Γ denote the semi-direct product of $\mathbb{Z}$ and H defined by

$$(f_1, n_1) \cdot (f_2, n_2) = (f_1 + n_1 \cdot f_2, n_1 + n_2).$$

It is easy to see that Γ is torsion free. For $k \in \mathbb{Z}$, we define $f^k \in H$ by $f^k(i) = 0$ if $k \neq i$ and $f^k(i) = 1$ if $k = i$. Then for any $k \in \mathbb{Z}$, $(0, 1)(f^{k+1}, 0) = (f^k, 0)(0, 1)$, i.e., $(0, 1)(f^{k+1}, 0)((0, 1)^{-1} = (f^k, 0)$. As $\{(0, f^k) : k \in \mathbb{Z}\}$ generates H as a $\mathbb{Z}$-module, this shows that Γ is generated by $(0, 1)$ and $(f^0, 0)$. In particular, Γ is finitely generated. For $k \geq 2$ let $\Gamma_k \subset \Gamma$ denote the set of all elements (f, n) such that $n = 0 \pmod{k}$ and

$$\sum_{i=-\infty}^{\infty} f(ki + j) = 0 (\mathrm{mod} k) \ \forall j = 0, 1, \ldots, k - 1.$$

It is easy to see that for each k, Γ_k is a normal subgroup of Γ and Γ/Γ_k is finite. Moreover for each non-zero $(f, n) \in \Gamma$ there exists k such that $(f, n) \notin \Gamma_k$. Hence Γ is residually finite.

Let X denote the compact abelian group $(\mathbb{Z}/2\mathbb{Z})^{\mathbb{Z}}$, equipped with the product topology and pointwise addition, and let $S : X \to X$ denote the shift map defined by $S(x)(i) = x(i + 1)$. We define an algebraic Γ-action α on X by $\alpha(f, n)(x) = S^n(x)$. It is easy to see that the shift action of $\mathbb{Z}$ on X is Noetherian. Since $\alpha|_H$ is trivial and the action of $\Gamma/H \cong \mathbb{Z}$ induced by α is the shift action on X, we deduce that α is also Noetherian.

Let $\pi : H \to X$ denote the homomorphism defined by $\pi(f)(i) = f(i) \ (mod\ 2)$. We define $c : \Gamma \to X$ by $c(f, n) = \pi(f)$. Then $c((f_1, n_1)(f_2, n_2)) = \pi(f_1 + n_1 \cdot f_2) = c(f_1, n_1) + S^{n_1}(\pi(f_2))$. Since

$$S^{n_1}(\pi(f_2)) = S^{n_1}(c(f_2, n_2)) = \alpha(f_1, n_1)(c(f_2, n_2)),$$

this shows that c is a 1-cocycle of α. We pick any $x \in X$ and define $c^{'} : \Gamma \to X$ by $c^{'}(\gamma) = \alpha(\gamma)(x) - x$. Let Λ be an arbitrary finite index subgroup of Γ. Since $\alpha(\gamma) = I$ for all $\gamma \in H$, it follows that $c^{'}(H \cap \Lambda) = \{0\}$. On the other hand, the restriction of c to H is a homomorphism with infinite image. As $H \cap \Lambda$ is a finite index subgroup of H, we deduce that $c(H \cap \Lambda)$ is also infinite. Hence $c|_\Lambda \neq c^{'}|_\Lambda$. As Λ and x are arbitrary, we conclude that c is not a virtual coboundary. We now apply Proposition 2.3. Let β denote the action corresponding to the cocycle c. Since α is Noetherian, so is β. Applying Proposition 2.1 we conclude that β does not have a dense set of periodic points.

3. Density of periodic orbits

In this section we concentrate on the case when Γ is polycyclic-by-finite. A countable group Γ is *polycyclic* if there exists a decreasing sequence of subgroups

$$\Gamma = \Gamma_n \supset \Gamma_{n-1} \supset \cdots \supset \Gamma_0 = \{0\}$$

such that for each i, Γ_i is a normal subgroup of Γ_{i+1} and Γ_{i+1}/Γ_i is cyclic. Any such series is called a *polycyclic series* of Γ. A group Γ is called *polycyclic-by-finite* if it contains a finite index subgroup that is polycyclic. If Γ is polycyclic-by-finite then every subgroup of Γ is finitely generated, and $\mathbb{Z}(\Gamma)$ is a Noetherian ring.

Let Γ be a polycyclic-by-finite group, $\Gamma_0 \subset \Gamma$ be a polycyclic subgroup of finite index and $\Gamma_0 = \Gamma_n \supset \Gamma_{n-1} \supset \cdots \supset \Gamma_0 = \{0\}$ be a polycyclic series of Γ_0. Then the number of i's such that Γ_{i+1}/Γ_i is infinite cyclic is independent of Γ_0 and the polycyclic series. This number is known as the *Hirsch number* of Γ. The following proposition summarizes some basic properties of this invariant (see [10]).

Proposition 3.1. *Let Γ be a polycyclic-by-finite group.*

(1) *$h(\Gamma) = 0$ if and only if Γ is finite.*

(2) *If* Γ_1 *is a subgroup of* Γ *then* $h(\Gamma_1) \leq h(\Gamma)$.
(3) *If* $\Gamma_1 \subset \Gamma$ *is normal then* $h(\Gamma) = h(\Gamma_1) + h(\Gamma/\Gamma_1)$.

We note that the previous proposition applies to finitely generated virtually nilpotent groups since they are polycyclic-by-finite. In the proof of Theorem 1.1 we will also use the following result about polycyclic-by-finite groups (see [8]):

Proposition 3.2. *Let* Γ *be a polycyclic-by-finite group and let* M *be a simple* $\mathbb{Z}[\Gamma]$*-module. Then* M *is finite.*

Our next lemma is a direct consequence of this result.

Lemma 3.3. *Let* Γ *be a polycyclic-by-finite group and let* (X, α) *be a Noetherian action of* Γ *on a non-trivial compact abelian group* X*. Then* X *admits a non-zero periodic point.*

Proof. Let $\mathcal{A}$ be the collection of all proper $\mathbb{Z}(\Gamma)$-submodules of $\widehat{X}$. Since α is Noetherian, $\mathcal{A}$ contains a maximal element M. It is easy to see that $N = \widehat{X}/M$ is a simple $\mathbb{Z}(\Gamma)$-module. By the previous proposition N is finite. Let $i : \widehat{N} \to X$ denote the dual of the projection map $\pi : \widehat{X} \to N$. Since π is surjective, i is injective. Hence $i(\widehat{N})$ is a non-trivial α-invariant finite subgroup of X. Clearly every non-zero point of $i(\widehat{N})$ is periodic. □

Lemma 3.4. *Let* Γ *be a polycyclic-by-finite group and let* α *be a Noetherian action of* Γ *on a zero-dimensional group* X *such that* $H^1_v(X, \alpha) \neq \{0\}$*. If* $\mathrm{Ker}(\alpha) = \{\gamma : \alpha(\gamma) = I\}$ *is infinite then there exists a polycyclic-by-finite group* Γ' *with* $h(\Gamma') < h(\Gamma)$*, and a Noetherian* Γ'*-action* α' *on* X *such that* $H^1_v(X, \alpha') \neq \{0\}$.

Proof. Let $\Lambda \subset \Gamma$ be a finite index subgroup, and let $c : \Lambda \to X$ be a virtual 1-cocycle of α that is not a virtual coboundary. Replacing Γ by Λ if necessary, we may assume that $\Lambda = \Gamma$. Since $\mathrm{Ker}(\alpha)$ acts trivially on X, it follows that the restriction of c to $\mathrm{Ker}(\alpha)$ is a homomorphism. Since Γ is polycyclic-by-finite, the subgroup $\mathrm{Ker}(\alpha)$ is finitely generated. This implies that $c(\mathrm{Ker}(\alpha))$ is also finitely generated. By Proposition 2.2 there exists $k \geq 2$ such that $kx = 0$ for

all $x \in X$. Hence $c(\mathrm{Ker}(\alpha))$ is a finite subgroup of X. We define $M = \{\gamma \in \mathrm{Ker}(\alpha) : c(\gamma) = 0\}$. Since $\mathrm{Ker}(\alpha)$ is finitely generated and $M \subset \mathrm{Ker}(\alpha)$ has finite index, M contains a finite index characteristic subgroup of $\mathrm{Ker}(\alpha)$, i.e., there exists a finite index subgroup $N \subset M$ such that $\theta(N) = N$ for all automorphism θ of $\mathrm{Ker}(\alpha)$. Since for any $g \in \Gamma$ the map $\gamma \mapsto g\gamma g^{-1}$ is an automorphism of $\mathrm{Ker}(\alpha)$, we deduce that N is a normal subgroup of Γ. We define $\Gamma' = \Gamma/N$. As N is a finite index subgroup of $\mathrm{Ker}(\alpha)$ and $\mathrm{Ker}(\alpha)$ is infinite, N is also infinite. In particular $h(N) > 0$. Hence

$$h(\Gamma/N) = h(\Gamma) - h(N) < h(\Gamma).$$

As $\alpha(\gamma) = I$ for all $\gamma \in N$, α induces a Noetherian Γ'-action α' on X. We define $c' : \Gamma' \to X$ by $c'(\gamma N) = c(\gamma)$. We note that for any $\gamma \in \Gamma$ and $n \in N$, $c(\gamma n) = c(\gamma) + \alpha(\gamma)(c(n)) = c(\gamma)$. Hence c' is a well defined virtual 1-cocycle of α'. Suppose there exist a finite index subgroup $\Lambda' \subset \Gamma'$ and $x \in X$ such that $c'(a) = \alpha'(a)(x) - x$ for all $a \in \Lambda'$. We define $\Lambda_1 = \pi^{-1}(\Lambda')$, where $\pi : \Gamma \to \Gamma'$ is the projection map. Then for any $\gamma \in \Lambda_1$,

$$c(\gamma) = c'(\pi(\gamma)) = \alpha'(\pi(\gamma))(x) - x = \alpha(\gamma)(x) - x.$$

Since this contradicts the fact that c is not a virtual coboundary, we conclude that c' is also not a virtual coboundary. Hence $H^1_v(X, \alpha') \neq 0$.□

Our next result shows that for Notherian actions of virtually nilpotent groups on zero-dimensional compact abelian groups, the virtual first cohomology group vanishes. In view of Theorem 2.5 and Lemma 3.3 this completes the proof of Theorem 1.1.

Theorem 3.5. *Let Γ be a finitely generated virtually nilpotent group, and let (X, α) be a Noetherian action of Γ on a zero-dimensional group X. Then $H^1_v(\alpha) = \{0\}$*

Proof. Let $\Lambda \subset \Gamma$ be a finite index subgroup, and let $c : \Lambda \to X$ be a virtual 1-cocycle of α. Suppose $\Gamma_0 \subset \Gamma$ is a nilpotent subgroup of finite index. Then c is equivalent to $c|_{\Gamma_0 \cap \Lambda}$. Hence, to show that c is

a virtual coboundary, without loss of generality we may assume that $\Lambda \subset \Gamma_0$.

We will use induction on $h(\Lambda)$, the Hirsch number of Λ. If $h(\Lambda)) = 0$ then Λ is finite, and hence c is a virtual coboundary. Suppose $h(\Lambda) \geq 1$. Then Λ is a finitely generated infinite nilpotent group. Hence the center of Λ is also finitely generated and infinite ([4], Proposition 2.8). We choose an element γ_0 in the center of Λ that has infinite order. For $m \geq 1$, we define $K_m = (\alpha(\gamma_0^m) - I)(X)$. Since $\alpha(\gamma_0)$ commutes with $\alpha(\gamma)$ for all $\gamma \in \Gamma$, it follows that each K_m is a closed α-invariant subgroup. As α is Noetherian, the collection $\{K_m : m \geq 1\}$ has a minimal element K. Clearly K is of the form $(\alpha(\gamma_1) - I)(X)$, where $\gamma_1 = \gamma_0^n$ for some $n \geq 1$. Let β denote the induced algebraic Γ-action on the quotient X/K. If P denotes the projection map from X to X/K then $P \circ c$ is virtual 1-cocycle of the action β. We note that the $\beta(\gamma) = I$ for all γ in the infinite cyclic subgroup generated by γ_1. By Lemma 3.4 and the induction hypothesis we deduce that $P \circ c$ is a virtual coboundary of β. We find a finite index subgroup $\Lambda_1 \subset \Lambda$ and $p \in X/K$ such that

$$P \circ c(\gamma) = \beta(\gamma)(p) - p \ \forall \gamma \in \Lambda_1.$$

We choose $q \in X$ such that $P(q) = p$. Let $c_1 : \Lambda_1 \to X$ denote the virtual 1-cocycle defined by $c_1 = c - c_q$. Since P is an equivariant map it follows that $P \circ c_1 = 0$, i.e., the image of c_1 is contained in $K = (\alpha(\gamma_1) - I)(X)$. We choose $l \geq 1$ such that $\gamma_1^l \in \Lambda_1$. From the minimality of K we deduce that $K = (\alpha(\gamma_2) - I)(X)$, where $\gamma_2 = \gamma_1^l$. We find $x \in X$ such that $\alpha(\gamma_2)(x) - x = c_1(\gamma_2)$. Let $c_2 : \Lambda_1 \to X$ be the virtual 1-cocycle defined by $c_2(\gamma) = c_1(\gamma) - c_x(\gamma)$. Then c_2 is cohomologous to c_1 and $c_2(\gamma_2) = 0$. Let $F \subset X$ denote the set of points that are fixed by $\alpha(\gamma_2)$. Since $\alpha(\gamma_2)$ commutes with $\alpha(\gamma)$ for all γ, F is a α-invariant closed subgroup. We note that for any $\gamma \in \Lambda_1$, $c_2(\gamma_2\gamma) = c_2(\gamma_2) + \alpha(\gamma_2)(c_2(\gamma)) = \alpha(\gamma_2)(c_2(\gamma))$. As γ_2 lies in the center of Γ we also have,

$$c_2(\gamma_2\gamma) = c_2(\gamma\gamma_2) = c_2(\gamma) + \alpha(\gamma)(c_2(\gamma_2)) = c_2(\gamma).$$

This shows that for all $\gamma \in \Lambda_1$ the element $c_2(\gamma)$ lies in F. Let α_F denote the restriction of α to F. Then c_2 can be viewed as a virtual 1-cocycle of α_F. Since $\alpha(\gamma)|_F = I$ for all γ in the infinite cyclic group generated by γ_2, from the previous lemma and the induction hypothesis we conclude that c_2 is a virtual coboundary. Since c is cohomologous to c_2 this completes the proof. □

We conclude this paper with the following question :

Question 3.6. Does Theorem 1.1 hold if Γ is polycyclic-by-finite ?

In view of Theorem 2.5 and Lemma 3.3 this is true if and only if Theorem 3.5 holds for polycyclic-by-finite groups.

References

[1] Lewis Bowen, *Entropy for expansive algebraic actions of residually finite groups*, Ergodic Theory Dynam. Systems 31 (2011), no. 3, 703-718.

[2] Nhan-Phu Chung, and Hanfeng Li, *Homoclinic groups, IE groups, and expansive algebraic actions*, Invent. Math. 199 (2015), no. 3, 805-858.

[3] Christopher Deninger, *Fuglede-Kadison determinants and entropy for actions of discrete amenable groups*, J. Amer. Math. Soc. 19 (2006), no. 3, 737-758.

[4] Meng-Che Ho, *Describing groups*, Proc. Amer. Math. Soc. 145 (2017), no. 5, 2223-2239.

[5] Bruce Kitchens and Klaus Schmidt, *Automorphisms of compact groups*, Ergodic theory dynam. systems 9 (1989), no. 4, 691-735.

[6] Hanfeng Li, *Compact group automorphisms, addition formulas and Fuglede-Kadison determinants*, Ann. of Math. (2) 176 (2012), no. 1, 303-347.

[7] Douglas Lind and Klaus Schmidt, *A survey of algebraic actions of the discrete Heisenberg group*, Russian Math. Surveys 70 (2015), no. 4, 657-714.

[8] J. E. Roseblade, *Group rings of polycyclic groups*, J. Pure Appl. Algebra 3 (1973), 307-328.

[9] Klaus Schmidt, *Dynamical systems of algebraic origin*, Birkhauser Verlag, 1995.

[10] B. A. F. Wehrfritz, *Group and ring theoretic properties of polycyclic groups*, Algebra and Applications, 10. Springer-Verlag London, Ltd., London, 2009.

School of Mathhematics, Tata Institute of Fundamental Research, Mumbai 400005, India

E-mail address: siddhart@math.tifr.res.in